

# t17\_euclid (TMVwUyCJDWzBcx- TYy7VyyuvYkAcifo1PCme)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k1\_euclid : \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k12\_euclid : \iota \Rightarrow \iota$  be given. Let  $k8\_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v3\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $m1\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v3\_membered : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Rightarrow ((v1\_xboole\_0 X0) \vee ((v2\_xxreal\_0 X1) \vee (v3\_xxreal\_0 X0)))))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (2)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(m2\_finseq\_2 X1 k1\_numbers (k1\_euclid X0)) \Rightarrow (\forall X2.(m2\_finseq\_2 X2 k1\_numbers (k1\_euclid X0)) \Rightarrow ((k12\_euclid (k8\_euclid X0 X1 X2) = k6\_numbers) \Leftrightarrow (X1 = X2)))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_2 X1 X0) \Rightarrow (\forall X2.(m2\_finseq\_2 X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (5)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (6)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0) \wedge ((v1\_xcmplx\_0 X0) \wedge ((v1\_xxreal\_0 X0) \wedge (v1\_xreal\_0 X0))) \quad (7)$$

Assume the following.

$$v3\_membered\ k1\_numbers \quad (8)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finseq\_1 X0) \wedge (v3\_valued\_0 X0)))) \Rightarrow (\neg v3\_xxreal\_0 (k12\_euclid X0)) \quad (9)$$

Assume the following.

$$v1\_xboole\_0\ k1\_xboole\_0 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_2 X1 X0) \Rightarrow (\forall X2.(m2\_finseq\_2 X2 X0 X1) \Rightarrow (m2\_finseq\_1 X2 X0)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge ((v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7\_ordinal1 X0) \wedge ((m1\_subset\_1 X1 (k1\_euclid X0)) \wedge (m1\_subset\_1 X2 (k1\_euclid X0)))) \Rightarrow (m2\_finseq\_2 (k8\_euclid X0 X1 X2) k1\_numbers (k1\_euclid X0)) \quad (14)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (m1\_finseq\_2 (k1\_euclid X0) k1\_numbers) \quad (15)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finseq\_1 X0) \wedge (v3\_valued\_0 X0)))) \Rightarrow (m1\_subset\_1 (k12\_euclid X0) k1\_numbers) \quad (16)$$

Assume the following.

$$\forall X0.((v1\_xreal\_0 X0) \wedge (v2\_xreal\_0 X0)) \Rightarrow ((\neg v1\_xboole\_0 X0) \wedge ((v1\_xreal\_0 X0) \wedge (\neg v3\_xreal\_0 X0))) \quad (17)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.(v3\_membered X1) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v3\_valued\_0 X2)) \quad (19)$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(m2\_finseq\_2 X1 k1\_numbers (k1\_euclid X0)) \Rightarrow (\forall X2.(m2\_finseq\_2 X2 k1\_numbers (k1\_euclid X0)) \Rightarrow (\neg(X1 \neq X2) \wedge (r1\_xreal\_0 (k12\_euclid (k8\_euclid X0 X1 X2)) k6\_numbers))))$$