

t17\_finseq\_1  
(TMThZxfaeFhAy5ofast4kfZ38exNfsJhx5J)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((k2\_finseq\_1 X0 = k2\_finseq\_1 X1) \Rightarrow (X0 = X1))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(v1\_relat\_1 X1) \Rightarrow ((r1\_tarski X0 (k9\_xtuple\_0 X1)) \Rightarrow (k9\_xtuple\_0 (k5\_relat\_1 X1 X0) = X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Leftrightarrow (r1\_tarski (k2\_finseq\_1 X0) (k2\_finseq\_1 X1)))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (6)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \quad (7)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (8)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (m2\_subset\_1 (k3\_finseq\_1 X0) k1\_numbers k5\_numbers) \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1. (m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow ((X1 = k3\_finseq\_1 \\ & X0) \Leftrightarrow (k2\_finseq\_1 X1 = k9\_xtuple\_0 X0))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0. (m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (11)$$

Assume the following.

$$\forall X0. (v1\_xboole\_0 X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (v1\_xboole\_0 X1)) \quad (12)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. (v7\_ordinal1 X0) \Rightarrow (\forall X1. ((v1\_relat\_1 X1) \wedge (( \\ & v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2))) \Rightarrow (((r1\_xreal\_0 X0 ( \\ & k3\_finseq\_1 X1)) \wedge (X2 = k5\_relat\_1 X1 (k2\_finseq\_1 X0))) \Rightarrow ((k3\_finseq\_1 \\ & X2 = X0) \wedge (k4\_finseq\_1 X2 = k2\_finseq\_1 X0)))))) \end{aligned}$$