

t17_graphsp (TMTHC- Ngh34G4aLaxmpHoTS3HchYqj7J2RQ)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_2 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_graphsp : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k13_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_graphsp : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & \forall X3.(X2 \neq X3) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X3 = k1_funct_1 \\ & \quad X0 X3)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X2 = X1)) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{3}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0.k3_finseq_2 X0 = k13_finseq_1 X0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X0 k1_numbers)\wedge(m1_subset_1 X2 k1_numbers))\Rightarrow(k1_graphsp X0 X1 X2 = k2_funct_7 X0 X1 X2) \quad (8)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (9)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Rightarrow(m1_subset_1 X2 X0)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Rightarrow(m2_finseq_1 X2 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (13)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (14)$$

Assume the following.

$$\forall X0.m1_finseq_2 (k3_finseq_2 X0) X0 \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X0 k1_numbers)\wedge(m1_subset_1 X2 k1_numbers))\Rightarrow(m2_finseq_1 (k1_graphsp X0 X1 X2) k1_numbers) \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (\forall X2.(m2_finseq_1 \\ & X2 k1_numbers) \Rightarrow (\forall X3.(m1_subset_1 X3 k1_numbers) \Rightarrow (k2_graphsp \\ & X0 X1 X2 X3 = k1_graphsp (k1_graphsp X2 X0 X1) X1 X3)))) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow (v5_relat_1 X1 X0) \quad (18)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v5_relat_1 X0 k1_numbers)) \Rightarrow ((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \quad (19)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (\forall X2.(m2_finseq_2 \\ & X2 k1_numbers (k3_finseq_2 k1_numbers)) \Rightarrow (\forall X3.(m1_subset_1 \\ & X3 k1_numbers) \Rightarrow ((X0 \in k4_finseq_1 X2) \Rightarrow ((X0 = X1) \vee (k1_seq_1 (k2_graphsp \\ & X0 X1 X2 X3) X0 = X1)))))) \end{aligned}$$