

t17\_polyform  
(TMag5QV3y56C7vNKuuCDr4FGE2piZv3ucZY)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & \quad (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ & \quad X1))) \Rightarrow (\forall X2.((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 \\ & \quad X2)))) \Rightarrow (k7\_finseq\_1 (k7\_finseq\_1 X0 X1) X2 = k7\_finseq\_1 X0 (k7\_finseq\_1 \\ & \quad X1 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ & \quad X1))) \Rightarrow ((k3\_finseq\_1 (k7\_finseq\_1 X1 (k9\_finseq\_1 X0)) = k2\_nat\_1 \\ & \quad (k3\_finseq\_1 X1) np\_1) \wedge ((k3\_finseq\_1 (k7\_finseq\_1 (k9\_finseq\_1 \\ & \quad X0) X1) = k2\_nat\_1 (k3\_finseq\_1 X1) np\_1) \wedge ((k1\_funct\_1 (k7\_finseq\_1 \\ & \quad X1 (k9\_finseq\_1 X0)) (k2\_nat\_1 (k3\_finseq\_1 X1) np\_1) = X0) \wedge (k1\_funct\_1 \\ & \quad (k7\_finseq\_1 (k9\_finseq\_1 X0) X1) np\_1 = X0)))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.v1\_finseq\_1 (k5\_finseq\_1 X0) \tag{4}$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k9\_finseq\_1 X0)) \wedge (v1\_funct\_1 (k9\_finseq\_1 X0)) \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.(((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 \\ X0)))\wedge((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))))\Rightarrow \\ ((v1\_relat\_1 (k7\_finseq\_1 X0 X1))\wedge((v1\_funct\_1 (k7\_finseq\_1 \\ X0 X1))\wedge(v1\_finseq\_1 (k7\_finseq\_1 X0 X1)))) \end{aligned} \quad (6)$$

**Theorem 1**

$$\begin{aligned} \forall X0.\forall X1.((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 \\ X1)))\Rightarrow(\forall X2.((v1\_relat\_1 X2)\wedge((v1\_funct\_1 X2)\wedge(v1\_finseq\_1 \\ X2)))\Rightarrow(k1\_funct\_1 (k7\_finseq\_1 (k7\_finseq\_1 (k9\_finseq\_1 X0) \\ X1) X2) np\_1 = X0)) \end{aligned}$$