

t17_procal_1

(TMT2uVfJTUodzcuL6BAy8c6udU2qdQUNoxe)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $k8_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_cqc_the1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow ((k8_cqc_lang X0 (k6_cqc_lang X0 X1) X2 \in \\ k4_cqc_the1 X0) \Rightarrow (k8_cqc_lang X0 (k6_cqc_lang X0 X2) X1 \in k4_cqc_the1 \\ X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X3.(m2_subset_1 X3 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (((k8_cqc_lang X0 X1 X2 \in k4_cqc_the1 X0) \wedge \\ (k8_cqc_lang X0 X2 X3 \in k4_cqc_the1 X0)) \Rightarrow (k8_cqc_lang X0 X1 X3 \in k4_cqc_the1 \\ X0)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (k8_cqc_lang X0 (k6_cqc_lang X0 (k8_cqc_lang \\ X0 X1 (k6_cqc_lang X0 X2))) (k7_cqc_lang X0 X1 X2) \in k4_cqc_the1 X0))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ X1\ X2)\ (k9_cqc_lang \\ X0\ (k6_cqc_lang\ X0\ X1)\ X2) \in k4_cqc_the1\ X0))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0\ X0) \wedge ((\neg v1_xboole_0\ X1) \wedge \\ (m1_subset_1\ X1\ (k1_zfmisc_1\ X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2\ X0\ X1) \Leftrightarrow (m1_subset_1\ X2\ X1)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(m1_qc_lang1\ X0) \Rightarrow (\neg v1_xboole_0\ (k3_cqc_lang\ X0)) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1\ X0) \wedge ((m1_subset_1 \\ X1\ (k3_cqc_lang\ X0)) \wedge (m1_subset_1\ X2\ (k3_cqc_lang\ X0)))) \Rightarrow (m2_subset_1 \\ (k9_cqc_lang\ X0\ X1\ X2)\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1\ X0) \wedge ((m1_subset_1 \\ X1\ (k3_cqc_lang\ X0)) \wedge (m1_subset_1\ X2\ (k3_cqc_lang\ X0)))) \Rightarrow (m2_subset_1 \\ (k8_cqc_lang\ X0\ X1\ X2)\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1\ X0) \wedge ((m1_subset_1 \\ X1\ (k3_cqc_lang\ X0)) \wedge (m1_subset_1\ X2\ (k3_cqc_lang\ X0)))) \Rightarrow (m2_subset_1 \\ (k7_cqc_lang\ X0\ X1\ X2)\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((m1_qc_lang1\ X0) \wedge (m1_subset_1\ X1\ (k3_cqc_lang \\ X0))) \Rightarrow (m2_subset_1\ (k6_cqc_lang\ X0\ X1)\ (k9_qc_lang1\ X0)\ (k3_cqc_lang \\ X0)) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.(m1_qc_lang1\ X0) \Rightarrow (m1_subset_1\ (k3_cqc_lang\ X0)\ (k1_zfmisc_1 \\ (k9_qc_lang1\ X0))) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1 \\ X0)) \Rightarrow (v1_xboole_0\ X1)) \quad (12)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (k8_cqc_lang\ X0\ (k6_cqc_lang\ X0\ (k7_cqc_lang \\ X0\ X1\ X2))\ (k9_cqc_lang\ X0\ (k6_cqc_lang\ X0\ X1)\ (k6_cqc_lang\ X0\ X2)) \in \\ k4_cqc_the1\ X0))) \end{aligned}$$