

t17_rinfsup1
(TMXnoW8Ze4TiQW8dfYoMkLRMFMr5qPHfvg2)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_partfun3 : \iota \Rightarrow o$ be given. Let $k1_valued_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $m2_valued_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (1)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((m1_subset_1 X0 k5_numbers) \wedge (v7_ordinal1 \\ & X1)) \Rightarrow (k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0))))))\wedge(v7_ordinal1 X2)))\Rightarrow(k1_valued_0 X0 X1 X2 = k9_nat_1 X1 X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (6)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (7)$$

Assume the following.

$$v3_membered k1_numbers \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0))))))\Rightarrow(\forall X2.(m2_valued_0 X2 X0 X1)\Rightarrow((v1_funct_1 X2)\wedge((v1_funct_2 X2 k5_numbers X0)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (10)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(m2_subset_1 (k2_nat_1 X0 X1) k1_numbers k5_numbers) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0))))))\wedge(v7_ordinal1 X2)))\Rightarrow(m2_valued_0 (k1_valued_0 X0 X1 X2) X0 X1) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((v4_partfun3 X0) \Leftrightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\ & (r1_xreal_0 k6_numbers (k1_seq_1 X0 X1)))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge (v1_partfun1 X0 k5_numbers)))) \Rightarrow (\forall X1.(v7_ordinal1 \\ & X1) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge ((v4_relat_1 X2 k5_numbers) \wedge \\ & ((v1_funct_1 X2) \wedge (v1_partfun1 X2 k5_numbers)))))) \Rightarrow ((X2 = k9_nat_1 \\ & X0 X1) \Leftrightarrow (\forall X3.(v7_ordinal1 X3) \Rightarrow (k1_funct_1 X2 X3 = k1_funct_1 \\ & X0 (k2_xcmplx_0 X3 X1)))))) \end{aligned} \quad (15)$$

Assume the following.

$$k1_xboole_0 = the (\lambda X0 : \iota.v1_xboole_0 X0) \quad (16)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(\neg v1_xboole_0 X1) \Rightarrow (\forall X2.(m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v1_funct_2 X2 X0 X1) \Rightarrow (\\ & v1_partfun1 X2 X0))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \quad (19)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \quad (20)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(v3_membered X1) \Rightarrow (\forall X2.(m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v3_valued_0 X2)) \end{aligned} \quad (21)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers k1_numbers) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow ((v4_partfun3 \\ & X1) \Rightarrow (v4_partfun3 (k1_valued_0 k1_numbers X1 X0))) \end{aligned}$$