

t17_subset_1 (TMEtwE98zYPrtFoxyBoiND5bvW78v59fJr6)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((r1_tarski X1 X2) \Leftrightarrow (r1_tarski \\ & (k3_subset_1 X0 X2) (k3_subset_1 X0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 X0 (k3_subset_1 X0 X1) = X1) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (m1_subset_1 (k3_subset_1 X0 X1) (k1_zfmisc_1 X0)) \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((r1_tarski (k3_subset_1 X0 \\ & X1) X2) \Rightarrow (r1_tarski (k3_subset_1 X0 X2) X1))) \end{aligned}$$