

t17\_waybel23  
(TMNokFnuts79bwDnCHpRfikqHYf7dMvZDPi)

October 27, 2020

Let  $v5\_orders\_2 : \iota \Rightarrow o$  be given. Let  $v2\_lattice3 : \iota \Rightarrow o$  be given. Let  $l1\_orders\_2 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_waybel23 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_yellow\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_domain\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $r2\_yellow\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg(X0 \in X1) \wedge ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X2)) \wedge (v1\_xboole\_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. ((v5\_orders\_2 X0) \wedge (l1\_orders\_2 X0)) \Rightarrow ((v2\_lattice3 X0) \Leftrightarrow (\forall X1. (m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (r2\_yellow\_0 X0 (k2\_tarski X1 X2)))))) \quad (3)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge (l1\_orders\_2 X0)) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow ((v1\_waybel23 X1 X0) \Leftrightarrow (\forall X2. (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3. (m1\_subset\_1 X3 (u1\_struct\_0 X0)) \Rightarrow (((X2 \in X1) \wedge ((X3 \in X1) \wedge (r2\_yellow\_0 X0 (k7\_domain\_1 (u1\_struct\_0 X0) X2 X3)))) \Rightarrow (k2\_yellow\_0 X0 (k7\_domain\_1 (u1\_struct\_0 X0) X2 X3) \in X1))))))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 X0) \wedge ((m1\_subset\_1 X1 X0) \wedge (m1\_subset\_1 X2 X0))) \Rightarrow (k7\_domain\_1 X0 X1 X2 = k2\_tarski X1 X2) \quad (5)$$

Assume the following.

$$\forall X0.(l1\_orders\_2 X0) \Rightarrow ((v2\_lattice3 X0) \Rightarrow (\neg v2\_struct\_0 X0)) \quad (6)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v5\_orders\_2 X0) \wedge ((v2\_lattice3 X0) \wedge (l1\_orders\_2 \\ & X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 \\ & X0))) \Rightarrow ((v1\_waybel23 X1 X0) \Leftrightarrow (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 \\ & X0)) \Rightarrow (\forall X3.(m1\_subset\_1 X3 (u1\_struct\_0 X0)) \Rightarrow (((X2 \in X1) \wedge \\ & (X3 \in X1)) \Rightarrow (k2\_yellow\_0 X0 (k7\_domain\_1 (u1\_struct\_0 X0) X2 X3) \in \\ & X1)))))) \end{aligned}$$