

t17\_xxreal\_0  
(TMQseLr9uCSVEvXmNqtsEXJmE8m1GSeV4Se)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (( \\ (r1\_xxreal\_0 X0 X1) \Rightarrow (k3\_xxreal\_0 X0 X1 = X0)) \wedge ((\neg r1\_xxreal\_0 X0 \\ X1) \Rightarrow (k3\_xxreal\_0 X0 X1 = X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow ( \\ (r1\_xxreal\_0 X0 X1) \vee (r1\_xxreal\_0 X1 X0)) \end{aligned} \tag{2}$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (r1\_xxreal\_0 \\ (k3\_xxreal\_0 X0 X1) X0)) \end{aligned}$$