

t181_relat_1
(TMPc8xkiSE9kiVekgLKC57abPD1S8Tx6CLm)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (r1_tarski X0 X1) \Rightarrow (r1_xboole_0 X0 (k4_xboole_0 X2 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k6_subset_1 X1 (k5_relat_1 X1 X0)) = k6_subset_1 (k9_xtuple_0 X1) X0) \quad (2)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. (r1_xboole_0 X1 (k9_xtuple_0 X0)) \Rightarrow (k5_relat_1 X0 X1 = k1_xboole_0)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X0) \Rightarrow (v1_relat_1 (k4_xboole_0 X0 X1)) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((r1_tarski X0 X1) \Rightarrow (k5_relat_1 (k6_subset_1 X2 (k5_relat_1 X2 X1)) X0 = k1_xboole_0))$$