

t185\_xreal\_1

(TMX81yoAo27cxijx7Rmprfhvz1PJLrVhzKq)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k7\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 k6\_numbers X1)) \Rightarrow (r1\_xxreal\_0 (k7\_xcmplx\_0 X0 X1) np\_1))) \quad (1)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 X1 X0)) \Rightarrow (r1\_xxreal\_0 (k7\_xcmplx\_0 X1 X0) np\_1)))$$