

t185_xxreal_1

(TMZxgZx9nKRWVYhcZH5mq2rPi979MHMPSgt)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k2_xxreal_1 X1 X2) \Leftrightarrow ((r1_xxreal_0 X1 X0) \wedge \\ & (\neg r1_xxreal_0 X2 X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X0) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v2_membered (k2_xxreal_1 X0 X1)) \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. \\ & (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (k2_xxreal_1 \\ X0 X1 = ReplSep (toset (\lambda X2 : \iota.m1_subset_1 X2 k7_numbers)) \\ (\lambda X2 : \iota.(r1_xxreal_0 X0 X2) \wedge (\neg r1_xxreal_0 X1 X2)) (\lambda X2 : \\ \iota.X2)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v2_membered X1) \Rightarrow ((\\ X0 = X1) \Leftrightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((X2 \in X0) \Leftrightarrow (X2 \in X1))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall X0.(v2_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow \\ (v1_xxreal_0 X1)) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ (v1_xxreal_0 X2) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k6_subset_1 (k2_xxreal_1 \\ X0 X2) (k2_xxreal_1 X0 X1) = k2_xxreal_1 X1 X2)))) \end{aligned}$$