

t186_xreal_1 (TMR-
byHLS1NFWhHxEBRg5Ap3i2zD2qgWeaPV)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (k7_xcmplx_0 X0 X0 = np_1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k7_xcmplx_0 X0 k6_numbers = k6_numbers) \quad (2)$$

Assume the following.

$$r1_xxreal_0 k6_numbers np_1 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\neg(r1_xxreal_0 X0 k6_numbers) \wedge (\neg r1_xxreal_0 \\ & (k7_xcmplx_0 X2 X0) (k7_xcmplx_0 X1 X0)) \wedge (r1_xxreal_0 X1 X2)))) \quad (4) \end{aligned}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (5)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 \\ & X0 k6_numbers) \wedge (r1_xxreal_0 X0 X1)) \Rightarrow (r1_xxreal_0 (k7_xcmplx_0 \\ & X1 X0) np_1))) \end{aligned}$$