

t188_xreal_1 (TMPuKbUAhGxct- LYC5LNoZRgtWHAoqpjt3jo)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 k6_numbers X0) \wedge ((\neg r1_xxreal_0 \\ & (k3_xcmplx_0 X2 X0) X1) \wedge (r1_xxreal_0 (k7_xcmplx_0 X1 X0) X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 np_1 X0 = X0) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg \\ & r1_xxreal_0 k6_numbers X0) \wedge ((\neg r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 \\ & (k7_xcmplx_0 X1 X0) np_1)))) \end{aligned}$$