

t18\_afinsq\_2  
(TMJsJXT4GF8NppbeWdFDEsvPnpUcUi2qTWF)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v5\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $k3\_afinsq\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_nat\_d : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k2\_afinsq\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((r1\_xxreal\_0 X1 X0) \Rightarrow (k7\_nat\_d X1 X0 = k6\_numbers))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (3)$$

Assume the following.

$$(m2\_subset\_1 np\_0 k1\_numbers k5\_numbers) \wedge ((m1\_subset\_1 np\_0 k5\_numbers) \wedge (m1\_subset\_1 np\_0 k1\_numbers)) \quad (4)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (5)$$

Assume the following.

$$r1\_xxreal\_0 np\_1 np\_1 \quad (6)$$

Assume the following.

$$r1\_xxreal\_0 np\_0 np\_1 \quad (7)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (8)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge ((v5\_ordinal1 \\ X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ (v7\_ordinal1 X2) \Rightarrow (k3\_afinsq\_2 X0 X1 X2 = k2\_afinsq\_2 (k5\_relat\_1 \\ X0 X2) (k7\_nat\_d X1 np\_1)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (11)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (( \\ v1\_funct\_1 X1) \wedge ((v5\_ordinal1 X1) \wedge (v1\_finset\_1 X1)))) \Rightarrow (k3\_afinsq\_2 \\ X1 k6\_numbers X0 = k3\_afinsq\_2 X1 np\_1 X0)) \end{aligned}$$