

# t18\_flang\_2 (TMcTxQ-GRhFy1BpqmVtjKCLWT6xwaHN2VXFa)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_catalan2 : \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (k2\_flang\_1 X0 \in k8\_flang\_1 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow ((k2\_flang\_1 X0 \in X1) \Rightarrow ((r1\_tarski X2 (k6\_flang\_1 X0 X2 X1)) \wedge (r1\_tarski X2 (k6\_flang\_1 X0 X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. k3\_catalan2 X0 = k8\_afinsq\_1 X0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (m1\_subset\_1 (k8\_flang\_1 X0 X1) (k1\_zfmisc\_1 (k3\_catalan2 X0))) \quad (4)$$

## Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow ((r1\_tarski X1 (k6\_flang\_1 X0 X1 (k8\_flang\_1 X0 X2))) \wedge (r1\_tarski X1 (k6\_flang\_1 X0 (k8\_flang\_1 X0 X2) X1)))) \end{aligned}$$