

# t18\_fvaluat1

(TMbS8oUmz9muA8KVRHMTU6xeHkteM24Vdry)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v6\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v3\_group\_1 : \iota \Rightarrow o$  be given. Let  $v5\_vectsp\_1 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v1\_realset2 : \iota \Rightarrow o$  be given. Let  $l6\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_fvaluat1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_fvaluat1 : \iota \Rightarrow o$  be given. Let  $k4\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_numbers : \iota$  be given. Let  $k1\_xxreal\_0 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_fvaluat1 : \iota \Rightarrow o$  be given. Let  $l2\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $l5\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $l2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $k4\_numbers : \iota$  be given. Let  $k6\_algstr\_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_algstr\_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1\_xboole\_0 X0) \wedge \\ & (((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))))) \wedge (m1\_subset\_1 X3 X0))) \Rightarrow (k3\_funct\_2 X0 \\ & X1 X2 X3 = k1\_funct\_1 X2 X3) \end{aligned} \quad (2)$$

Assume the following.

$$\neg v1\_xreal\_0 k1\_xxreal\_0 \quad (3)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge (l1\_struct\_0 X0)) \Rightarrow (\neg v1\_xboole\_0 (u1\_struct\_0 X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(l6\_algstr\_0 X0) \Rightarrow (\forall X1.(m1\_fvaluat1 X1 X0) \Rightarrow \\ ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 (u1\_struct\_0 X0) k7\_numbers) \wedge \\ ((v2\_fvaluat1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ (u1\_struct\_0 X0) k7\_numbers)))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(l6\_algstr\_0 X0) \Rightarrow ((l2\_algstr\_0 X0) \wedge (l5\_algstr\_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.(l2\_algstr\_0 X0) \Rightarrow ((l2\_struct\_0 X0) \wedge (l1\_algstr\_0 X0)) \quad (7)$$

Assume the following.

$$\forall X0.(l1\_algstr\_0 X0) \Rightarrow (l1\_struct\_0 X0) \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.(l6\_algstr\_0 X0) \Rightarrow ((v3\_fvaluat1 X0) \Rightarrow (\forall X1.( \\ (v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 (u1\_struct\_0 X0) k7\_numbers) \wedge \\ ((v2\_fvaluat1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ (u1\_struct\_0 X0) k7\_numbers)))))) \Rightarrow ((m1\_fvaluat1 X1 X0) \Leftrightarrow ((k1\_funct\_1 \\ X1 (k4\_struct\_0 X0) = k1\_xxreal\_0) \wedge ((\forall X2.(m1\_subset\_1 \\ X2 (u1\_struct\_0 X0) \Rightarrow ((X2 \neq k4\_struct\_0 X0) \Rightarrow (k1\_funct\_1 X1 X2 \in \\ k4\_numbers))) \wedge ((\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0) \Rightarrow \\ (\forall X3.(m1\_subset\_1 X3 (u1\_struct\_0 X0) \Rightarrow (k1\_funct\_1 X1 \\ (k6\_algstr\_0 X0 X2 X3) = k1\_xxreal\_3 (k1\_funct\_1 X1 X2) (k1\_funct\_1 \\ X1 X3)))) \wedge ((\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0) \Rightarrow ((r1\_xxreal\_0 \\ k6\_numbers (k1\_funct\_1 X1 X2) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k1\_funct\_1 \\ X1 (k1\_algstr\_0 X0 (k5\_struct\_0 X0) X2)))))) \wedge (\exists X2.(m1\_subset\_1 \\ X2 (u1\_struct\_0 X0) \wedge ((k1\_funct\_1 X1 X2 \neq k6\_numbers) \wedge (k1\_funct\_1 \\ X1 X2 \neq k1\_xxreal\_0)))))))))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.(v1\_int\_1 X0) \Rightarrow (v1\_xreal\_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_numbers) \Rightarrow (v1\_int\_1 X0) \quad (11)$$

**Theorem 1**

$$\begin{aligned} \forall X0.((\neg v2\_struct\_0 X0) \wedge ((\neg v6\_struct\_0 X0) \wedge ((v13\_algstr\_0 \\ X0) \wedge ((v3\_group\_1 X0) \wedge ((v5\_vectsp\_1 X0) \wedge ((v2\_rlvect\_1 X0) \wedge \\ (v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v1\_realset2 X0) \wedge (l6\_algstr\_0 \\ X0)))))))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow \\ (\forall X2.(m1\_fvaluat1 X2 X0) \Rightarrow (\neg (v3\_fvaluat1 X0) \wedge ((X1 \neq k4\_struct\_0 \\ X0) \wedge (k3\_funct\_2 (u1\_struct\_0 X0) k7\_numbers X2 X1 = k1\_xxreal\_0)))))) \end{aligned}$$