

t18_trees_2 (TMY-
DAqU82QAB4x1QmqWJSBrQdBd78aB2u6n)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_trees_1 : \iota \Rightarrow o$ be given. Let $m1_trees_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_trees_2 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $k1_trees_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_card_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge (v1_trees_1 X0)) \Rightarrow (\neg (v1_trees_2 \\ & X0) \wedge (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (\exists X2. \\ & (m1_trees_1 X2 X0) \wedge (\forall X3.(v1_finset_1 X3) \Rightarrow (\neg (X3 = k1_trees_2 \\ & X0 X2) \wedge (r1_xreal_0 (k5_card_1 X3) X1)))))) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge (v1_trees_1 X0)) \Rightarrow (\forall X1. \\ & (m1_trees_1 X1 X0) \Rightarrow ((v1_trees_2 X0) \Rightarrow (v1_finset_1 (k1_trees_2 \\ & X0 X1)))) \end{aligned}$$