

t18\_wsierp\_1  
(TMH43zb2CqxdE2vvYTdJA5D6EZhQty74Uby)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 (k6\_xcmplx\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 \\ & (k2\_xcmplx\_0 X0 X2) X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((r1\_xxreal\_0 (k2\_xcmplx\_0 X0 X1) X2) \Rightarrow ((r1\_xxreal\_0 \\ & X0 X2) \wedge (r1\_xxreal\_0 X1 X2)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_xreal\_0 X0) \tag{3}$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((r1\_xxreal\_0 X0 (k6\_xcmplx\_0 X1 X2)) \Rightarrow ((r1\_xxreal\_0 \\ & X0 X1) \wedge (r1\_xxreal\_0 X2 X1)))))) \end{aligned}$$