

# t18\_xboolean (TM- Sra2L9AvBwMcsTkTZj2k3ugmT45WcXLFs)

October 27, 2020

Let  $v1\_xboolean : \iota \Rightarrow o$  be given. Let  $k4\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xboolean : \iota \Rightarrow \iota$  be given. Let  $k6\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k4\_xboolean X0 (k5\_xboolean X0 X1) = k5\_xboolean X0 (k4\_xboolean X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (v1\_xboolean (k3\_xboolean X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k6\_xboolean X0 X1 = k5\_xboolean (k3\_xboolean X0) X1)) \quad (3)$$

## Theorem 1

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k4\_xboolean (k3\_xboolean X0) (k6\_xboolean X0 X1) = k5\_xboolean (k3\_xboolean X0) (k4\_xboolean (k3\_xboolean X0) X1)))$$