

t18_xxreal_2
(TMGitU9SkYbYLPvjgfc34FTyaQgzGy4gDcp)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_xxreal_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k3_xxreal_1 X1 X2) \Leftrightarrow ((\neg r1_xxreal_0 X0 X1) \wedge \\ & (r1_xxreal_0 X0 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v2_membered (k3_xxreal_1 X0 X1)) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ & m2_xxreal_2 X1 X0) \Leftrightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((X2 \in X0) \Rightarrow (r1_xxreal_0 \\ & X1 X2)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0) \tag{4}$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (m2_xxreal_2 X0 (k3_xxreal_1 X0 X1)))$$