

t190\_xreal\_1

(TMK1tP6eGLJ1HTWHgPr9rApWxywtSnPtFB9)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k7\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\neg(\neg r1\_xxreal\_0 k6\_numbers X0) \wedge ((\neg r1\_xxreal\_0 \\ & X2 (k3\_xcmplx\_0 X1 X0)) \wedge (r1\_xxreal\_0 X1 (k7\_xcmplx\_0 X2 X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k3\_xcmplx\_0 np\_1 X0 = X0) \quad (2)$$

Assume the following.

$$m1\_subset\_1 k1\_xboole\_0 k4\_ordinal1 \quad (3)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow ( \\ & r1\_xxreal\_0 X0 X2)))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow ((X0 \neq k6\_numbers) \Rightarrow (k7\_xcmplx\_0 X0 X0 = np\_1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (v1\_xreal\_0 (k7\_xcmplx\_0 X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_xreal\_0 X0) \quad (10)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\neg(r1\_xxreal\_0 X0 k6\_numbers) \wedge ((\neg r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 np\_1 (k7\_xcmplx\_0 X0 X1))))))$$