

t193_xcplx_1 (TMZTcJS-
RGm9hEhd2NKY2Fwc9eNaLa8AsNyg)

October 27, 2020

Let $v1_xcplx_0 : \iota \Rightarrow o$ be given. Let $k4_xcplx_0 : \iota \Rightarrow \iota$ be given. Let $k7_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (k7_xcplx_0 X0 np_1 = X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (k7_xcplx_0 X0 (k4_xcplx_0 X1) = k4_xcplx_0 (k7_xcplx_0 X0 X1))) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xcplx_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (k4_xcplx_0 X0 = k7_xcplx_0 X0 (k4_xcplx_0 np_1))$$