

t196_xreal_1 (TMNbfmPB-
mqVPV94eHQwb1h1hDbXZdMyJ8YZ)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 (k4_xcmplx_0 X1)) \Rightarrow (r1_xxreal_0 X1 (k4_xcmplx_0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 X1 (k4_xcmplx_0 X0))) \Rightarrow (r1_xxreal_0 (k4_xcmplx_0 np_1) (k7_xcmplx_0 X1 X0)))) \quad (2)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 X0 (k4_xcmplx_0 X1))) \Rightarrow (r1_xxreal_0 (k4_xcmplx_0 np_1) (k7_xcmplx_0 X1 X0))))$$