

t196_xxreal_1 (TMX-
ofb1e66yJAqmZNLpNZvGb1ArdmKpx2Zk)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. k2_xboole_0 X0 X1 = k5_xboole_0 X0 (k4_xboole_0 X1 X0) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. (v1_xxreal_0 X2) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (r1_tarski (k2_xxreal_1 X1 X2) (k4_xxreal_1 X0 X2)))))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. k4_xboole_0 (k4_xboole_0 X0 X1) X2 = k4_xboole_0 X0 (k2_xboole_0 X1 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (k4_xboole_0 X0 X1 = k1_xboole_0) \Leftrightarrow (r1_tarski X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow (r1_xxreal_0 X0 X2)))) \quad (5)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k1_xxreal_1 X1 X0 = k1_xboole_0))) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k6_subset_1 (k1_xxreal_1 \\ & X2 X1) (k3_xxreal_1 X0 X1) = k1_xxreal_1 X2 X0)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ & r1_xxreal_0 X0 X1) \Rightarrow (k6_subset_1 (k1_xxreal_1 X0 X1) (k1_tarski \\ & X1) = k2_xxreal_1 X0 X1))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ & \neg r1_xxreal_0 X1 X0) \Rightarrow (k3_xxreal_1 X0 X1 = k2_xboole_0 (k4_xxreal_1 \\ & X0 X1) (k1_tarski X1)))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (10)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ & (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (13)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k6_subset_1 (k2_xxreal_1 \\ & X2 X1) (k4_xxreal_1 X0 X1) = k1_xxreal_1 X2 X0)))) \end{aligned}$$