

t19\_complex1  
(TMLd1Mmo5HbMb91GBAvUYL6ZQGTvikQP1UR)

October 27, 2020

Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k3\_complex1 : \iota \Rightarrow \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_complex1 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_complex1 : \iota$  be given. Let  $k2\_numbers : \iota$  be given. Let  $k11\_complex1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (v1\_xreal\_0 X0) \Rightarrow (\forall X1. (v1\_xreal\_0 X1) \Rightarrow ((k3\_complex1 \\ (k2\_xcmplx\_0 X0 (k3\_xcmplx\_0 X1 k7\_complex1)) = X0) \wedge (k4\_complex1 \\ (k2\_xcmplx\_0 X0 (k3\_xcmplx\_0 X1 k7\_complex1)) = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((m1\_subset\_1 X0 k2\_numbers) \wedge (m1\_subset\_1 X1 k2\_numbers)) \Rightarrow (k11\_complex1 X0 X1 = k6\_xcmplx\_0 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 X1)) \Rightarrow (m1\_subset\_1 (k9\_real\_1 X0 X1) k1\_numbers) \quad (4)$$

Assume the following.

$$\forall X0. (v1\_xcmplx\_0 X0) \Rightarrow (m1\_subset\_1 (k4\_complex1 X0) k1\_numbers) \quad (5)$$

Assume the following.

$$\forall X0. (v1\_xcmplx\_0 X0) \Rightarrow (m1\_subset\_1 (k3\_complex1 X0) k1\_numbers) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_subset\_1 X0 k2\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 \\ X1 k2\_numbers) \Rightarrow (k11\_complex1 X0 X1 = k2\_xcmplx\_0 (k9\_real\_1 (k3\_complex1 \\ X0) (k3\_complex1 X1)) (k3\_xcmplx\_0 (k9\_real\_1 (k4\_complex1 X0) \\ (k4\_complex1 X1)) k7\_complex1))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Leftrightarrow (X0 \in k2\_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (9)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmplx\_0 X1) \Rightarrow (( \\ k3\_complex1 (k6\_xcmplx\_0 X0 X1) = k9\_real\_1 (k3\_complex1 X0) (k3\_complex1 \\ X1)) \wedge (k4\_complex1 (k6\_xcmplx\_0 X0 X1) = k9\_real\_1 (k4\_complex1 \\ X0) (k4\_complex1 X1)))) \end{aligned}$$