

t19_finseq_1

(TMYjBwxcXagBCHhnuRrjPcBQinkMGh9xs25)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X2 X0))\Rightarrow(k8_funcop_1 X0 X1 X2 = k2_funcop_1 X1 X2) \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (10)$$

Assume the following.

$$\forall X0.\exists X1.m1_subset_1 X1 X0 \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X2 X0))\Rightarrow((v1_funct_1 (k8_funcop_1 X0 X1 X2))\wedge((v1_funct_2 (k8_funcop_1 X0 X1 X2) X1 X0)\wedge(m1_subset_1 (k8_funcop_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X1 X0)))))) \quad (12)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1)))\Rightarrow((m1_finseq_1 X1 X0)\Leftrightarrow(r1_tarski (k10_xtuple_0 X1) X0)) \quad (14)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers)\Rightarrow((X1 = k3_finseq_1 X0)\Leftrightarrow(k2_finseq_1 X1 = k9_xtuple_0 X0))) \quad (15)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow((v1_finseq_1 X0)\Leftrightarrow(\exists X1.(v7_ordinal1 X1)\wedge(k9_xtuple_0 X0 = k2_finseq_1 X1))) \quad (16)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Leftrightarrow(X0 \in k4_ordinal1) \quad (17)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (19)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow (\exists X2.(m2_finseq_1 X2 X1) \wedge (k3_finseq_1 X2 = X0)))$$