

t19_frechet
(TMYQfpg2PiiMjjFSEi7LhV82G85RwJ5SpoZ)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_xboole_0 X0 X1) \Leftrightarrow (k4_xboole_0 X0 X1 = X0) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (\neg X0 \in X1) \Rightarrow (r1_xboole_0 (k1_tarski X0) X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (r1_tarski X2 X0) \Rightarrow \\ & (k8_relat_1 (k1_funct_4 (k6_partfun1 X0) (k7_funcop_1 X1 X3)) \\ & (k6_subset_1 X2 (k1_tarski X3)) = k7_subset_1 X2 (k6_subset_1 X2 \\ & X1) (k1_tarski X3)) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(r1_xboole_0 X0 X1)\Rightarrow(r1_xboole_0 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(m1_subset_1 (k7_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.m1_subset_1 (k6_subset_1 X0 X1) (k1_zfmisc_1 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_xboole_0 X1)) \quad (13)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.\forall X3.(r1_tarski X2 X0)\Rightarrow((X3 \in X0)\vee(k8_relat_1 (k1_funct_4 (k6_partfun1 X0) (k7_funcop_1 X1 X3)) (k6_subset_1 X2 (k1_tarski X3)) = k6_subset_1 X2 X1))$$