

t19_glib_002

(TMF5XsW6jjMecv7tJomWJDwe8chLgUYT1HA)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $k2_glib_002 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m3_glib_001 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_glib_001 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \neg (X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1. (m3_glib_001 \\ & X1 X0) \Rightarrow (\forall X2. \forall X3. \forall X4. \forall X5. ((v2_glib_001 \\ & X1 X0) \wedge ((r1_glib_001 X0 X2 X3 X1) \wedge (r2_glib_000 X0 X3 X5 X4))) \Rightarrow ((\\ & v2_glib_001 (k10_glib_001 X0 X1 X4) X0) \wedge (r1_glib_001 X0 X2 X5 (k10_glib_001 \\ & X0 X1 X4)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge \\ & ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \wedge (m1_subset_1 \\ & X1 (k6_glib_000 X0)) \Rightarrow ((\neg v1_xboole_0 (k2_glib_002 X0 X1)) \wedge (m1_subset_1 \\ & (k2_glib_002 X0 X1) (k1_zfmisc_1 (k6_glib_000 X0)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1_relat_1 X0) \wedge ((v4_relat_1 \\ & X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 \\ & X0)))))) \wedge (m3_glib_001 X1 X0) \Rightarrow (m3_glib_001 (k10_glib_001 X0 X1 \\ & X2) X0) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\
& X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\
& X1 (k6_glib_000 X0)) \Rightarrow (\forall X2.((\neg v1_xboole_0 X2) \wedge (m1_subset_1 \\
& X2 (k1_zfmisc_1 (k6_glib_000 X0)))) \Rightarrow ((X2 = k2_glib_002 X0 X1) \Leftrightarrow \\
& (\forall X3.(X3 \in X2) \Leftrightarrow (\exists X4.((v2_glib_001 X4 X0) \wedge (m3_glib_001 \\
& X4 X0)) \wedge (r1_glib_001 X0 X1 X3 X4)))))) \quad (5)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\
& X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1. \forall X2. \\
& \forall X3. \forall X4.(m1_subset_1 X4 (k6_glib_000 X0)) \Rightarrow (((X1 \in \\
& k2_glib_002 X0 X4) \wedge (r2_glib_000 X0 X1 X3 X2)) \Rightarrow (X3 \in k2_glib_002 \\
& X0 X4))
\end{aligned}$$