

t19_hausdorf
(TMFC96ZSGAhiiD7s78SNjmoC3tvBRPqvupH)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v6_metric_1 : \iota \Rightarrow o$ be given. Let $v7_metric_1 : \iota \Rightarrow o$ be given. Let $v8_metric_1 : \iota \Rightarrow o$ be given. Let $v9_metric_1 : \iota \Rightarrow o$ be given. Let $l1_metric_1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k3_pcomps_1 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_weierstr : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_metric_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_pre_topc : \iota \Rightarrow \iota$ be given. Let $k2_pcomps_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v6_metric_1 X0) \wedge ((v7_metric_1 X0) \wedge ((v8_metric_1 X0) \wedge ((v9_metric_1 X0) \wedge (l1_metric_1 X0)))))) \Rightarrow \\ & (\forall X1. ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 (k3_pcomps_1 X0))))) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 X0) \Rightarrow (\forall X3. (m1_subset_1 X3 (u1_struct_0 X0) \Rightarrow ((X3 \in X1) \Rightarrow (r1_xxreal_0 (k1_funct_1 (k6_weierstr X0 X1) X2) (k4_metric_1 X0 X2 X3))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. (l1_metric_1 X0) \Rightarrow ((u1_struct_0 X0 = u1_struct_0 (k3_pcomps_1 X0)) \wedge (u1_pre_topc (k3_pcomps_1 X0) = k2_pcomps_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Leftrightarrow (\forall X1. \neg X1 \in X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((v8_metric_1 X0) \wedge (l1_metric_1 X0)) \wedge ((m1_subset_1 X1 (u1_struct_0 X0)) \wedge (m1_subset_1 X2 (u1_struct_0 X0)))) \Rightarrow (k4_metric_1 X0 X1 X2 = k4_metric_1 X0 X2 X1) \quad (5)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v6_metric_1 X0) \wedge ((v7_metric_1 \\ & X0) \wedge ((v8_metric_1 X0) \wedge ((v9_metric_1 X0) \wedge (l1_metric_1 X0)))))) \Rightarrow \\ & (\forall X1.((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (u1_struct_0 (k3_pcomps_1 X0))))) \Rightarrow (\forall X2.(m1_subset_1 \\ & X2 (u1_struct_0 X0)) \Rightarrow (\exists X3.(m1_subset_1 X3 (u1_struct_0 \\ & X0)) \wedge ((X3 \in X1) \wedge (r1_xxreal_0 (k1_funct_1 (k6_weierstr X0 X1) X2) \\ & (k4_metric_1 X0 X3 X2))))) \end{aligned}$$