

t19_prob_2

(TMQxFtdoyz1hfujiq6xummYHxBUJzuTPQav)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_prob_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X1) \wedge ((v1_prob_1 X1 X0) \wedge \\ & ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. (m1_prob_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (((\neg v1_xboole_0 X1) \wedge ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0)))))) \wedge ((m1_subset_1 X2 X1) \wedge (m1_subset_1 X3 X1))) \Rightarrow \\ & (k5_prob_1 X0 X1 X2 X3 = k3_xboole_0 X2 X3) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (v1_xreal_0 (k1_funct_1 X0 X1)) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge \\ & ((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.(m2_prob_1 X2 X0 X1)\Rightarrow((v1_funct_1 \\ & X2)\wedge((v1_funct_2 X2 X1 k1_numbers)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X1 k1_numbers)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0))\Rightarrow(m1_subset_1 (k1_seq_1 X0 X1) k1_numbers) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge \\ & ((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.(m2_prob_1 X2 X0 X1)\Rightarrow(\forall X3. \\ & (m1_prob_1 X3 X0 X1)\Rightarrow(\forall X4.(m1_prob_1 X4 X0 X1)\Rightarrow((r1_prob_2 \\ & X0 X1 X2 X3 X4)\Leftrightarrow(k1_seq_1 X2 (k5_prob_1 X0 X1 X3 X4) = k8_real_1 (k1_seq_1 \\ & X2 X3) (k1_seq_1 X2 X4)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(k8_real_1 X0 X1 = k8_real_1 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (12)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge \\ & ((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.(m1_prob_1 X2 X0 X1)\Rightarrow(\forall X3. \\ & (m1_prob_1 X3 X0 X1)\Rightarrow(\forall X4.(m2_prob_1 X4 X0 X1)\Rightarrow((r1_prob_2 \\ & X0 X1 X4 X2 X3)\Rightarrow(r1_prob_2 X0 X1 X4 X3 X2)))))) \end{aligned}$$