

t19_rfunct_2

(TMbr25bXhTXiFb3dTX9FLAk1Efm5k4Ryn1D)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v5_xxreal_2 : \iota \Rightarrow o$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_seq_4 : \iota \Rightarrow \iota$ be given. Let $k5_seq_4 : \iota \Rightarrow \iota$ be given. Let $v3_funct_1 : \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ ((v5_xxreal_2 (k2_relset_1 k1_numbers X1)) \wedge (k4_seq_4 (k2_relset_1 \\ k1_numbers X1) = k5_seq_4 (k2_relset_1 k1_numbers X1))) \Rightarrow (v3_funct_1 \\ X1))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k10_xtuple_0 (k5_relat_1 X1 X0) = k7_relat_1 X1 X0) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (\\ k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k7_relset_1 X0 X1 X2 X3 = k7_relat_1 \\ X2 X3) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (\\ k2_relset_1 X0 X1 = k10_xtuple_0 X1) \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow(k2_partfun1 \\ & X0 X1 X2 X3 = k5_relat_1 X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 \\ & (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) \\ & (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.((v1_funct_1 \\ & X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k1_numbers))))\Rightarrow \\ & (((v5_xxreal_2 (k7_relset_1 X1 k1_numbers X2 X0))\wedge(k4_seq_4 (\\ & k7_relset_1 X1 k1_numbers X2 X0) = k5_seq_4 (k7_relset_1 X1 k1_numbers \\ & X2 X0)))\Rightarrow(v3_funct_1 (k2_partfun1 X1 k1_numbers X2 X0)))) \end{aligned}$$