

t19_sprect_1

(TMUnKo2SYfpTQfR3qZyiarW2agS91FK9FBF)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v2_compts_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k11_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k10_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $v1_sppol_1 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k7_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k6_pscomp_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow ((k19_euclid X0 \\ & X1 = k19_euclid X2 X3) \Rightarrow ((X0 = X2) \wedge (X1 = X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow \\ & ((v1_sppol_1 X0) \Leftrightarrow (k9_pscomp_1 X0 = k7_pscomp_1 X0)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k9_pscomp_1 X0) k1_numbers) \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k7_pscomp_1 X0) k1_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k6_pscomp_1 X0) k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (k11_pscomp_1 X0 = k19_euclid (k6_pscomp_1 X0) (k7_pscomp_1 X0)) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (k10_pscomp_1 X0 = k19_euclid (k6_pscomp_1 X0) (k9_pscomp_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow ((k11_pscomp_1 X0 = k10_pscomp_1 X0) \Rightarrow (v1_sppol_1 X0))$$