

t19_substut1
(TMYvZtmXTrA9tRhfi5qhEHso35jU6wFq6kb)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k16_substut1 : \iota \Rightarrow \iota$ be given. Let $k19_substut1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k29_substut1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k21_substut1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v6_substut1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_qc_lang1 X0) \wedge ((m1_subset_1 X1 (k16_substut1 X0)) \wedge (m1_subset_1 X2 (k16_substut1 X0)))) \Rightarrow (m1_subset_1 (k21_substut1 X0 X1 X2) (k16_substut1 X0)) \quad (1)$$

Assume the following.

$$\forall X0. (m1_qc_lang1 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k16_substut1 X0)) \Rightarrow ((v6_substut1 X1 X0) \Rightarrow (\forall X2. (m1_subset_1 X2 (k16_substut1 X0)) \Rightarrow ((X2 = k29_substut1 X0 X1) \Leftrightarrow (\exists X3. (m1_subset_1 X3 (k16_substut1 X0)) \wedge ((X1 = k21_substut1 X0 X3 X2) \wedge (k19_substut1 X0 X3 = k19_substut1 X0 X2)))))))) \quad (2)$$

Assume the following.

$$\forall X0. (m1_qc_lang1 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k16_substut1 X0)) \Rightarrow ((v6_substut1 X1 X0) \Leftrightarrow (\exists X2. (m1_subset_1 X2 (k16_substut1 X0)) \wedge (\exists X3. (m1_subset_1 X3 (k16_substut1 X0)) \wedge ((X1 = k21_substut1 X0 X2 X3) \wedge (k19_substut1 X0 X2 = k19_substut1 X0 X3)))))) \quad (3)$$

Theorem 1

$$\forall X0. (m1_qc_lang1 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k16_substut1 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 (k16_substut1 X0)) \Rightarrow ((k19_substut1 X0 X1 = k19_substut1 X0 X2) \Rightarrow (k29_substut1 X0 (k21_substut1 X0 X1 X2) = X2))))$$