

t19_tdlat_2

(TMKVyy3dRjdetYKTf6NE128sMuWVZbtw6Ln)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k1_tdlat_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \neg (X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0. \exists X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (v1_xboole_0 X1) \quad (5)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (6)$$

Assume the following.

$$\forall X0. \exists X1. m1_subset_1 X1 X0 \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. ((l1_pre_topc X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow (m1_subset_1 (k1_tops_1 X0 X1) (k1_zfmisc_1 (u1_struct_0 X0))) \quad (8)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v2_pre_topc\ X0) \wedge (l1_pre_topc\ X0)) \Rightarrow (\forall X1. \\
& (m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))) \Rightarrow \\
& (\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0 \\
& X0)))) \Rightarrow ((X2 = k1_tdlat_2\ X0\ X1) \Leftrightarrow (\forall X3.(m1_subset_1\ X3\ (k1_zfmisc_1 \\
& (u1_struct_0\ X0)))) \Rightarrow ((X3 \in X2) \Leftrightarrow (\exists X4.(m1_subset_1\ X4\ (k1_zfmisc_1 \\
& (u1_struct_0\ X0)))) \wedge ((X3 = k1_tops_1\ X0\ X4) \wedge (X4 \in X1))))))
\end{aligned} \tag{9}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v2_pre_topc\ X0) \wedge (l1_pre_topc\ X0)) \Rightarrow (\forall X1. \\
& (m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))) \Rightarrow \\
& ((X1 = k1_xboole_0) \Leftrightarrow (k1_tdlat_2\ X0\ X1 = k1_xboole_0))
\end{aligned}$$