

t19_zf_lang
(TMZxEi6PtdnAX7rzwoxfLWdFJeFGSryqZ7Z)

October 27, 2020

Let $v1_zf_lang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v3_zf_lang : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zf_lang : \iota$ be given. Let $k4_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k5_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k5_numbers k1_zf_lang) \Rightarrow (\forall X1. \\ & (m2_subset_1 X1 k5_numbers k1_zf_lang) \Rightarrow ((k1_funct_1 (k4_zf_lang \\ & X0 X1) np_1 = k6_numbers) \wedge (k1_funct_1 (k5_zf_lang X0 X1) np_1 = \\ & np_1))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow ((\\ & v3_zf_lang X0) \Leftrightarrow (\exists X1.(m2_subset_1 X1 k5_numbers k1_zf_lang) \wedge \\ & (\exists X2.(m2_subset_1 X2 k5_numbers k1_zf_lang) \wedge (X0 = k5_zf_lang \\ & X1 X2)))) \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow ((\\ & v3_zf_lang X0) \Rightarrow (k1_funct_1 X0 np_1 = np_1)) \end{aligned}$$