

t1\_aofa\_i00

(TMHK1VknWJePseZoeWf2odmnqC8xaF94Grm)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k16\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k7\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. k1\_funct\_1 (k16\_funcop\_1 X0 X1) X0 = X1 \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2. \\ & ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((X0 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 \\ & (k1\_funct\_4 X2 X1) X0 = k1\_funct\_1 X1 X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (k9\_xtuple\_0 (k2\_funcop\_1 X0 X1) = X0) \wedge (r1\_tarski (k10\_xtuple\_0 (k2\_funcop\_1 X0 X1)) (k1\_tarski X1)) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2. \\ & ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((\neg X0 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 \\ & (k1\_funct\_4 X2 X1) X0 = k1\_funct\_1 X2 X0))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k7\_funcop\_1 X0 X1 = k2\_funcop\_1 X0 X1 \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 (k2\_funcop\_1 X0 X1)) \wedge (v1\_funct\_1 (k2\_funcop\_1 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(((v1\_relat\_1 X0)\wedge(v1\_funct\_1 X0))\wedge((v1\_relat\_1 X1)\wedge(v1\_funct\_1 X1)))\Rightarrow((v1\_relat\_1 (k1\_funct\_4 X0 X1))\wedge(v1\_funct\_1 (k1\_funct\_4 X0 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k16\_funcop\_1 X0 X1 = k7\_funcop\_1 (k1\_tarski X0) X1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.k2\_funcop\_1 X0 X1 = k2\_zfmisc\_1 X0 (k1\_tarski X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1\_tarski X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow(X2 = X0)) \quad (10)$$

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.\forall X5. \neg((X3\neq X4)\wedge((X4\neq X5)\wedge((X5\neq X3)\wedge(\forall X6.((v1\_relat\_1 X6)\wedge(v1\_funct\_1 X6))\Rightarrow(\neg(k1\_funct\_1 X6 X3 = X0)\wedge((k1\_funct\_1 X6 X4 = X1)\wedge(k1\_funct\_1 X6 X5 = X2)))))))$$