

t1_bcialg_2

(TMPa6MPG2uXQNp6VQ53dJWmEFimCwXNgHB4)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v3_bcialg_1 : \iota \Rightarrow o$ be given. Let $v4_bcialg_1 : \iota \Rightarrow o$ be given. Let $v5_bcialg_1 : \iota \Rightarrow o$ be given. Let $v7_bcialg_1 : \iota \Rightarrow o$ be given. Let $l2_bcialg_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_bcialg_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_bcialg_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$m1_subset_1 \ k1_xboole_0 \ k4_ordinal1 \tag{1}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{2}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (((\neg v2_struct_0 \\ & X0) \wedge ((v3_bcialg_1 X0) \wedge ((v4_bcialg_1 X0) \wedge ((v5_bcialg_1 X0) \wedge \\ & ((v7_bcialg_1 X0) \wedge (l2_bcialg_1 X0)))))) \wedge ((m1_subset_1 X1 (u1_struct_0 \\ & X0)) \wedge ((m1_subset_1 X2 (u1_struct_0 X0)) \wedge (m1_subset_1 X3 k5_numbers)))) \Rightarrow \\ & (m1_subset_1 (k1_bcialg_2 X0 X1 X2 X3) (u1_struct_0 X0)) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_bialg_1 X0) \wedge ((v4_bialg_1 \\
& X0) \wedge ((v5_bialg_1 X0) \wedge ((v7_bialg_1 X0) \wedge (l2_bialg_1 X0)))))) \Rightarrow \\
& (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 \\
& X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 k5_numbers) \Rightarrow \\
& (\forall X4.(m1_subset_1 X4 (u1_struct_0 X0)) \Rightarrow ((X4 = k1_bialg_2 \\
& X0 X1 X2 X3) \Leftrightarrow (\exists X5.((v1_funct_1 X5) \wedge ((v1_funct_2 X5 k5_numbers \\
& (u1_struct_0 X0)) \wedge (m1_subset_1 X5 (k1_zfmisc_1 (k2_zfmisc_1 \\
& k5_numbers (u1_struct_0 X0)))))) \wedge ((X4 = k3_funct_2 k5_numbers \\
& (u1_struct_0 X0) X5 X3) \wedge ((k3_funct_2 k5_numbers (u1_struct_0 \\
& X0) X5 k6_numbers = X1) \wedge (\forall X6.(m1_subset_1 X6 k5_numbers) \Rightarrow \\
& ((\neg r1_xreal_0 X3 X6) \Rightarrow (k3_funct_2 k5_numbers (u1_struct_0 X0) \\
& X5 (k2_nat_1 X6 np_1) = k1_bialg_1 X0 (k3_funct_2 k5_numbers (\\
& u1_struct_0 X0) X5 X6) X2))))))))))
\end{aligned} \tag{5}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_bialg_1 X0) \wedge ((v4_bialg_1 \\
& X0) \wedge ((v5_bialg_1 X0) \wedge ((v7_bialg_1 X0) \wedge (l2_bialg_1 X0)))))) \Rightarrow \\
& (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 \\
& X2 (u1_struct_0 X0)) \Rightarrow (k1_bialg_2 X0 X1 X2 k6_numbers = X1)))
\end{aligned}$$