

t1_chord (TMUUPCrtQz- ZQMheUigXx7NzbQiAWtUpof4q)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k4_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg(k6_numbers \neq X0) \wedge (r1_xxreal_0 X0 k6_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k4_nat_d X0 X0 = k6_numbers) \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (m2_subset_1 (k6_xcmplx_0 X1 np_1) k1_numbers k5_numbers))) \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((\neg r1_xxreal_0 np_1 X0) \Rightarrow (X0 = k6_numbers)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (8)$$

Assume the following.

$$(\neg v1_xboole_0 \ k4_ordinal1) \wedge (v3_ordinal1 \ k4_ordinal1) \quad (9)$$

Assume the following.

$$v1_xboole_0 \ k1_xboole_0 \quad (10)$$

Assume the following.

$$m1_subset_1 \ k5_numbers \ (k1_zfmisc_1 \ k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. ((v7_ordinal1 \ X0) \wedge (v7_ordinal1 \ X1)) \Rightarrow (m1_subset_1 \ (k4_nat_d \ X0 \ X1) \ k5_numbers) \quad (12)$$

Assume the following.

$$\forall X0. (m1_subset_1 \ X0 \ k4_ordinal1) \Rightarrow (v7_ordinal1 \ X0) \quad (13)$$

Assume the following.

$$\forall X0. (v1_xboole_0 \ X0) \Rightarrow (\forall X1. (m1_subset_1 \ X1 \ (k1_zfmisc_1 \ X0)) \Rightarrow (v1_xboole_0 \ X1)) \quad (14)$$

Theorem 1

$$\forall X0. ((\neg v1_xboole_0 \ X0) \wedge (v7_ordinal1 \ X0)) \Rightarrow ((v7_ordinal1 \ (k6_xcmplx_0 \ X0 \ np_1)) \wedge (r1_xxreal_0 \ np_1 \ X0))$$