

t1_closure2 (TMXYK- gRYvCjk8p3uCdh1W6HU33HzazqHtgM)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (k1_funct_1 (k4_relat_1 X1) X0 = X0) \quad (2)$$

Assume the following.

$$\forall X0. k6_partfun1 X0 = k4_relat_1 X0 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1_xboole_0 X0) \wedge \\ & (((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1)))))) \wedge (m1_subset_1 X3 X0))) \Rightarrow (k3_funct_2 X0 \\ & X1 X2 X3 = k1_funct_1 X2 X3) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. (v1_relat_1 (k4_relat_1 X0)) \wedge ((v4_relat_1 (k4_relat_1 X0) X0) \wedge ((v1_funct_1 (k4_relat_1 X0)) \wedge (v1_partfun1 (k4_relat_1 X0) X0))) \quad (5)$$

Assume the following.

$$\forall X0. (v1_partfun1 (k6_partfun1 X0) X0) \wedge (m1_subset_1 (k6_partfun1 X0) (k1_zfmisc_1 (k2_zfmisc_1 X0 X0))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v1_partfun1 X2 X0)\Rightarrow(v1_funct_2 X2 X0 X1)) \quad (7)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(\forall X2.(m1_subset_1 X2 X0)\Rightarrow((r1_tarski X1 X2)\Rightarrow(r1_tarski (k3_funct_2 X0 X0 (k6_partfun1 X0) X1) (k3_funct_2 X0 X0 (k6_partfun1 X0) X2))))))$$