

t1_dist_1

(TMJ5MZ9YETGbDSzqCBAABxkM6zRK1YVE4wj)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k8_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((k7_relset_1 X0 X1 X2 X0 = k2_relset_1 X1 \\ & X2) \wedge (k8_relset_1 X0 X1 X2 X1 = k1_relset_1 X0 X2)) \end{aligned} \tag{1}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & k1_relset_1 X0 X1 = k9_xtuple_0 X1) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ & (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))))) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \tag{6}$$

Theorem 1

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow(k9_xtuple_0 X1 = k8_reset_1 \\ k5_numbers X0 X1 X0)$$