

t1\_fintopo5  
(TMaCsyh5f5VX42k6fzDr3myaMyMzYk34Wfe)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k7\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_funct\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. ((v2\_funct\_1 X0) \wedge (r1\_tarski X1 (k9\_xtuple\_0 X0))) \Rightarrow (k7\_relat\_1 (k2\_funct\_1 X0) (k7\_relat\_1 X0 X1) = X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k7\_relset\_1 X0 X1 X2 X3 = k7\_relat\_1 X2 X3) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow (((X1 \neq k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 X2 X0 \\ & X1) \Leftrightarrow (X0 = k1\_relset\_1 X0 X2))) \wedge ((X1 = k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 \\ & X2 X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0)))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \end{aligned} \quad (8)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.\forall X1.(\neg v1\_xboole\_0 X1) \Rightarrow (\forall X2.((v1\_funct\_1 \\ & X2) \wedge ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1)))))) \Rightarrow (\forall X3.(m1\_subset\_1 X3 (k1\_zfmisc\_1 X0)) \Rightarrow ((v2\_funct\_1 \\ & X2) \Rightarrow (k7\_relat\_1 (k2\_funct\_1 X2) (k7\_relset\_1 X0 X1 X2 X3) = X3)))) \end{aligned}$$