

t1_hilbert1 (TMRURhwkqYSdVJFhaftiwvMg- WaKDkVJb1vf)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_hilbert1 : \iota$ be given. Let $k2_hilbert1 : \iota$ be given. Let $k5_hilbert1 : \iota \Rightarrow \iota$ be given. Let $v6_hilbert1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_hilbert1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_hilbert1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 k1_hilbert1)) \Rightarrow (m1_subset_1 (k5_hilbert1 X0) (k1_zfmisc_1 k1_hilbert1)) \quad (1)$$

Assume the following.

$$m1_subset_1 k2_hilbert1 k1_hilbert1 \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 k1_hilbert1)) \Rightarrow (\forall X1. \\ (m1_subset_1 X1 (k1_zfmisc_1 k1_hilbert1)) \Rightarrow ((X1 = k5_hilbert1 \\ X0) \Leftrightarrow (\forall X2.(m1_subset_1 X2 k1_hilbert1) \Rightarrow ((X2 \in X1) \Leftrightarrow (\forall X3. \\ (m1_subset_1 X3 (k1_zfmisc_1 k1_hilbert1)) \Rightarrow (((v6_hilbert1 X3) \wedge \\ (r1_tarski X0 X3)) \Rightarrow (X2 \in X3)))))))) \quad (3) \end{aligned}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 k1_hilbert1)) \Rightarrow (((v6_hilbert1 \\ X0) \Leftrightarrow ((k2_hilbert1 \in X0) \wedge (\forall X1.(m1_subset_1 X1 k1_hilbert1) \Rightarrow \\ (\forall X2.(m1_subset_1 X2 k1_hilbert1) \Rightarrow (\forall X3.(m1_subset_1 \\ X3 k1_hilbert1) \Rightarrow ((k3_hilbert1 X1 (k3_hilbert1 X2 X1) \in X0) \wedge ((k3_hilbert1 \\ (k3_hilbert1 X1 (k3_hilbert1 X2 X3)) (k3_hilbert1 (k3_hilbert1 \\ X1 X2) (k3_hilbert1 X1 X3)) \in X0) \wedge ((k3_hilbert1 (k4_hilbert1 X1 \\ X2) X1 \in X0) \wedge ((k3_hilbert1 (k4_hilbert1 X1 X2) X2 \in X0) \wedge ((k3_hilbert1 \\ X1 (k3_hilbert1 X2 (k4_hilbert1 X1 X2)) \in X0) \wedge (((X1 \in X0) \wedge (k3_hilbert1 \\ X1 X2 \in X0)) \Rightarrow (X2 \in X0)))))))))))) \quad (4) \end{aligned}$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 k1_hilbert1)) \Rightarrow (k2_hilbert1 \in k5_hilbert1 X0)$$