

t1_jordan16 (TMUTPwM- LeYYHrHAA4g3rYiGMGusZw9UJPBr)

October 27, 2020

Let $v1_topreal2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k9_jordan6 : \iota \Rightarrow \iota$ be given. Let $k8_jordan6 : \iota \Rightarrow \iota$ be given. Let $r1_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k22_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k1_jordan5c : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_jordan6 : \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k8_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k2_jordan5c : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\
 & \quad np_2)))) \Rightarrow ((v1_topreal2 X0) \Rightarrow ((r1_topreal1 (k15_euclid np_2) \\
 & (k18_pscomp_1 X0) (k22_pscomp_1 X0) (k8_jordan6 X0)) \wedge ((r1_topreal1 \\
 & (k15_euclid np_2) (k22_pscomp_1 X0) (k18_pscomp_1 X0) (k8_jordan6 \\
 & X0)) \wedge ((r1_topreal1 (k15_euclid np_2) (k22_pscomp_1 X0) (k18_pscomp_1 \\
 & X0) (k9_jordan6 X0)) \wedge ((r1_topreal1 (k15_euclid np_2) (k18_pscomp_1 \\
 & X0) (k22_pscomp_1 X0) (k9_jordan6 X0)) \wedge ((k9_subset_1 (u1_struct_0 \\
 & (k15_euclid np_2)) (k8_jordan6 X0) (k9_jordan6 X0) = k2_tarski \\
 & (k18_pscomp_1 X0) (k22_pscomp_1 X0)) \wedge ((k4_subset_1 (u1_struct_0 \\
 & (k15_euclid np_2)) (k8_jordan6 X0) (k9_jordan6 X0) = X0) \wedge (\neg r1_xxreal_0 \\
 & (k18_euclid (k1_jordan5c (k8_jordan6 X0) (k6_jordan6 (k10_real_1 \\
 & (k7_real_1 (k6_pscomp_1 X0) (k8_pscomp_1 X0)) np_2)) (k18_pscomp_1 \\
 & X0) (k22_pscomp_1 X0))) (k18_euclid (k2_jordan5c (k9_jordan6 \\
 & X0) (k6_jordan6 (k10_real_1 (k7_real_1 (k6_pscomp_1 X0) (k8_pscomp_1 \\
 & X0)) np_2)) (k22_pscomp_1 X0) (k18_pscomp_1 X0)))))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 (k1_zfmisc_1 (u1_struct_0 (k15_euclid X0)))) \Rightarrow (\forall X2. \\ (m1_subset_1 X2 (u1_struct_0 (k15_euclid X0))) \Rightarrow (\forall X3.(\\ m1_subset_1 X3 (u1_struct_0 (k15_euclid X0))) \Rightarrow (\neg(r1_topreal1 \\ (k15_euclid X0) X2 X3 X1) \wedge (\forall X4.(m1_subset_1 X4 (u1_struct_0 \\ (k15_euclid X0))) \Rightarrow (\neg(X4 \in X1) \wedge ((X4 \neq X2) \wedge (X4 \neq X3)))))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_subset_1 X1 (k1_zfmisc_1 \\ X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (k4_subset_1 X0 X1 X1 = \\ X1) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X0 = X0 \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ np_2)))) \Rightarrow ((\neg v1_xboole_0 (k9_jordan6 X0)) \wedge (m1_subset_1 (k9_jordan6 \\ X0) (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ np_2)))) \Rightarrow (m1_subset_1 (k22_pscomp_1 X0) (u1_struct_0 (k15_euclid \\ np_2))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ np_2)))) \Rightarrow (m1_subset_1 (k18_pscomp_1 X0) (u1_struct_0 (k15_euclid \\ np_2))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(X2 = k2_tarski X0 X1) \Leftrightarrow (\forall X3. \\ (X3 \in X2) \Leftrightarrow ((X3 = X0) \vee (X3 = X1))) \end{aligned} \quad (10)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_topreal2 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 \\ (k15_euclid np_2)))))) \Rightarrow (k9_jordan6 X0 \neq k8_jordan6 X0) \end{aligned}$$