

t1_jordan9

(TMYHz7JH7FV3VW2YxPKqgMdvDGvHBqBhDte)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v2_connsp_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r3_connsp_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_xboole_0 X1 X2)) \Rightarrow (r1_xboole_0 X0 X2) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc X0))) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X3. (m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X4. (m1_subset_1 X4 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (((v2_connsp_1 X2 X0) \wedge (r3_connsp_1 X0 X4 X3) \wedge (r1_tarski X1 X3) \wedge (r1_tarski X2 X4)))) \Rightarrow ((r1_xboole_0 X1 X2) \vee (r1_tarski X2 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X3. \\ & (m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\neg (r3_connsp_1 X0 X3 X1) \wedge (r3_connsp_1 X0 X3 X2) \wedge ((X1 \neq X2) \wedge (\neg r1_xboole_0 X1 X2)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r1_xboole_0 X0 X1) \Rightarrow (r1_xboole_0 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \quad (5)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (\forall X4.(m1_subset_1 X4 (k1_zfmisc_1 (u1_struct_0 \\ & X0)))) \Rightarrow (((v2_connsp_1 X1 X0) \wedge ((r3_connsp_1 X0 X4 X2) \wedge ((r3_connsp_1 \\ & X0 X4 X3) \wedge (r1_tarSKI X1 X4)))) \Rightarrow ((r1_xboole_0 X1 X2) \vee ((r1_xboole_0 \\ & X1 X3) \vee (X2 = X3))))))))) \end{aligned}$$