

# t1\_matrix\_1

(TMaTY2jvNhwFGXeRzSB6owbr7zdGuZHWEay)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_matrix\_1 : \iota \Rightarrow o$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((X1 = k9\_finseq\_1 X0) \Leftrightarrow ((k3\_finseq\_1 X1 = np\_1) \wedge (k10\_xtuple\_0 X1 = k1\_tarski X0))) \quad (1)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (2)$$

Assume the following.

$$\forall X0. k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge (m1\_subset\_1 X1 X0)) \Rightarrow (k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (5)$$

Assume the following.

$$\forall X0. v1\_finseq\_1 (k5\_finseq\_1 X0) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k5\_finseq\_1 X0)) \wedge (v1\_funct\_1 (k5\_finseq\_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1\_tarski X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ ((v1\_matrix\_1 X0) \Leftrightarrow (\exists X1.(v7\_ordinal1 X1) \wedge (\forall X2. \\ \neg(X2 \in k10\_xtuple\_0 X0) \wedge (\forall X3.((v1\_relat\_1 X3) \wedge ((v1\_funct\_1 \\ X3) \wedge (v1\_finseq\_1 X3)))) \Rightarrow (\neg(X3 = X2) \wedge (k3\_finseq\_1 X3 = X1)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (10)$$

**Theorem 1**

$$\forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow (v1\_matrix\_1 (k9\_finseq\_1 (k12\_finseq\_1 X0 X1))))$$