

t1_mesfunc5 (TMTgAyGUjZdNeRp- mViXBS2ZKERK3DqB72QL)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $k3_extreal1 : \iota \Rightarrow \iota$ be given. Let $k4_supinf_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k2_xxreal_3 : \iota \Rightarrow \iota$ be given. Let $k3_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_supinf_2 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((k2_xxreal_3 (k3_xxreal_3 X0 X1) = k1_xxreal_3 (k2_xxreal_3 X0 X1) \wedge (k2_xxreal_3 (k3_xxreal_3 X0 X1) = k3_xxreal_3 X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k7_numbers) \Rightarrow (k3_extreal1 X0 = k3_extreal1 (k2_supinf_2 X0)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k7_numbers) \wedge (m1_subset_1 X1 k7_numbers)) \Rightarrow (k4_supinf_2 X0 X1 = k3_xxreal_3 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k7_numbers) \Rightarrow (k2_supinf_2 X0 = k2_xxreal_3 X0) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k7_numbers) \wedge (m1_subset_1 X1 k7_numbers)) \Rightarrow (m1_subset_1 (k4_supinf_2 X0 X1) k7_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k7_numbers) \Rightarrow (v1_xxreal_0 X0) \quad (6)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k7_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k7_numbers) \Rightarrow (k3_extreal1 (k4_supinf_2 X0 X1) = k3_extreal1 (k4_supinf_2 X1 X0)))$$