

## t1\_modelc\_2

(TMWjK94zLn67Lo2PNCWNWSBBfagVE2Dr8d1)

October 27, 2020

Let  $v1\_modelc\_2 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k9\_modelc\_2 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_modelc\_2 : \iota \Rightarrow \iota$  be given. Let  $k3\_modelc\_2 : \iota \Rightarrow \iota$  be given. Let  $k4\_modelc\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_modelc\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_modelc\_2 : \iota \Rightarrow \iota$  be given. Let  $k7\_modelc\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_modelc\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$k5\_numbers = k4\_ordinal1 \tag{1}$$

Assume the following.

$$\neg v1\_xboole\_0 \ k9\_modelc\_2 \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow ((X0 = k9\_modelc\_2) \Leftrightarrow ((\forall X1. \\
& (X1 \in X0) \Rightarrow (m2\_finseq\_1 X1 k5\_numbers)) \wedge ((\forall X1.(v7\_ordinal1 \\
& X1) \Rightarrow (k2\_modelc\_2 X1 \in X0)) \wedge ((\forall X1.(m2\_finseq\_1 X1 k5\_numbers) \Rightarrow \\
& ((X1 \in X0) \Rightarrow (k3\_modelc\_2 X1 \in X0))) \wedge ((\forall X1.(m2\_finseq\_1 X1 \\
& k5\_numbers) \Rightarrow (\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow (((X1 \in \\
& X0) \wedge (X2 \in X0)) \Rightarrow (k4\_modelc\_2 X1 X2 \in X0)))) \wedge ((\forall X1.(m2\_finseq\_1 \\
& X1 k5\_numbers) \Rightarrow (\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow (((X1 \in \\
& X0) \wedge (X2 \in X0)) \Rightarrow (k5\_modelc\_2 X1 X2 \in X0)))) \wedge ((\forall X1.(m2\_finseq\_1 \\
& X1 k5\_numbers) \Rightarrow ((X1 \in X0) \Rightarrow (k6\_modelc\_2 X1 \in X0))) \wedge ((\forall X1. \\
& (m2\_finseq\_1 X1 k5\_numbers) \Rightarrow (\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow \\
& (((X1 \in X0) \wedge (X2 \in X0)) \Rightarrow (k7\_modelc\_2 X1 X2 \in X0)))) \wedge ((\forall X1. \\
& (m2\_finseq\_1 X1 k5\_numbers) \Rightarrow (\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow \\
& (((X1 \in X0) \wedge (X2 \in X0)) \Rightarrow (k8\_modelc\_2 X1 X2 \in X0)))) \wedge (\forall X1.( \\
& \neg v1\_xboole\_0 X1) \Rightarrow (((\forall X2.(X2 \in X1) \Rightarrow (m2\_finseq\_1 X2 k5\_numbers)) \wedge \\
& ((\forall X2.(v7\_ordinal1 X2) \Rightarrow (k2\_modelc\_2 X2 \in X1)) \wedge ((\forall X2. \\
& (m2\_finseq\_1 X2 k5\_numbers) \Rightarrow ((X2 \in X1) \Rightarrow (k3\_modelc\_2 X2 \in X1))) \wedge \\
& ((\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow (\forall X3.(m2\_finseq\_1 \\
& X3 k5\_numbers) \Rightarrow (((X2 \in X1) \wedge (X3 \in X1)) \Rightarrow (k4\_modelc\_2 X2 X3 \in X1)))) \wedge \\
& ((\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow (\forall X3.(m2\_finseq\_1 \\
& X3 k5\_numbers) \Rightarrow (((X2 \in X1) \wedge (X3 \in X1)) \Rightarrow (k5\_modelc\_2 X2 X3 \in X1)))) \wedge \\
& ((\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow ((X2 \in X1) \Rightarrow (k6\_modelc\_2 \\
& X2 \in X1))) \wedge ((\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow (\forall X3. \\
& (m2\_finseq\_1 X3 k5\_numbers) \Rightarrow (((X2 \in X1) \wedge (X3 \in X1)) \Rightarrow (k7\_modelc\_2 \\
& X2 X3 \in X1)))) \wedge (\forall X2.(m2\_finseq\_1 X2 k5\_numbers) \Rightarrow (\forall X3. \\
& (m2\_finseq\_1 X3 k5\_numbers) \Rightarrow (((X2 \in X1) \wedge (X3 \in X1)) \Rightarrow (k8\_modelc\_2 \\
& X2 X3 \in X1))))))))) \Rightarrow (r1\_tarski X0 X1))))))))) \Rightarrow (3)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \Rightarrow ((m1\_subset\_1 X1 X0) \Leftrightarrow \\
& (X1 \in X0))) \wedge ((v1\_xboole\_0 X0) \Rightarrow ((m1\_subset\_1 X1 X0) \Leftrightarrow (v1\_xboole\_0 \\
& X1))) \Rightarrow (4)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(m2\_finseq\_1 X0 k5\_numbers) \Rightarrow ((v1\_modelc\_2 X0) \Leftrightarrow (m1\_subset\_1 \\
& X0 k9\_modelc\_2)) \Rightarrow (5)
\end{aligned}$$

**Theorem 1**

$$\begin{aligned}
& \forall X0.((v1\_modelc\_2 X0) \wedge (m2\_finseq\_1 X0 k5\_numbers)) \Leftrightarrow ( \\
& X0 \in k9\_modelc\_2)
\end{aligned}$$