

t1_pencil_4

(TMTF4GedWp59WSffCW6HANP6giZk76w8UoM)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_3 : \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\neg \\ & (r1_xxreal_0 X0 (k1_nat_1 X1 np_1)) \wedge (\neg r1_xxreal_0 X0 X1) \wedge (X0 \neq \\ & k1_nat_1 X1 np_1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_xxreal_0 (k2_xcmplx_0 \\ & X0 X2) (k2_xcmplx_0 X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k2_xcmplx_0 np_2 np_1 = np_3 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v7_ordinal1 X0) \wedge (m1_subset_1 X1 k5_numbers)) \Rightarrow \\ & (k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xreal_0\ X0) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers)\Rightarrow(v1_xreal_0\ X0) \quad (8)$$

Theorem 1

$$\begin{aligned} &\forall X0.(v7_ordinal1\ X0)\Rightarrow(\forall X1.(v7_ordinal1\ X1)\Rightarrow(\neg \\ &(\neg r1_xxreal_0\ X1\ X0)\wedge((r1_xxreal_0\ np_3\ X1)\wedge((r1_xxreal_0\ X1 \\ &(k1_nat_1\ X0\ np_1))\wedge(\neg r1_xxreal_0\ np_2\ X0)))) \end{aligned}$$