

t1_polynom3 (TMTKYP- NMp2D5mTX8aL4SENU562QheXfW4ww)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $k4_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (1)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v3_rlvect_1 \\ & X0) \wedge ((v4_rlvect_1 X0) \wedge (l2_algstr_0 X0)))) \Rightarrow (\forall X1. (m2_finseq_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow ((\forall X2. (m1_subset_1 X2 k5_numbers) \Rightarrow \\ & ((X2 \in k4_finseq_1 X1) \Rightarrow (k1_funct_1 X1 X2 = k4_struct_0 X0)) \Rightarrow (k4_rlvect_1 \\ & X0 X1 = k4_struct_0 X0)))) \end{aligned} \quad (3)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (4)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v3_rlvect_1 \\ & X0) \wedge ((v4_rlvect_1 X0) \wedge (l2_algstr_0 X0)))))) \Rightarrow (\forall X1.(m2_finseq_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow ((\forall X2.(m2_subset_1 X2 k1_numbers \\ & k5_numbers) \Rightarrow ((X2 \in k4_finseq_1 X1) \Rightarrow (k1_funct_1 X1 X2 = k4_struct_0 \\ & X0))) \Rightarrow (k4_rlvect_1 X0 X1 = k4_struct_0 X0))) \end{aligned}$$